

Distribution-Free Testing Algorithms for Monomials with a Sublinear Number of Queries

Elya Dolev
School of Electrical Engineering
Tel-Aviv University
Ramat Aviv, ISRAEL
elyadolev@msn.com

Dana Ron*
School of Electrical Engineering
Tel-Aviv University
Ramat Aviv, ISRAEL
danar@eng.tau.ac.il

June 19, 2010

Abstract

We consider the problem of distribution-free testing of the class of monotone monomials and the class of monomials over n variables. While there are very efficient algorithms for testing a variety of functions classes when the underlying distribution is uniform, designing distribution-free algorithms (which must work under any arbitrary and unknown distribution), tends to be a more challenging task. When the underlying distribution is uniform, Parnas et al. (*SIAM Journal on Discrete Math*, 2002) give an algorithm for testing (monotone) monomials whose query complexity does not depends on n , and whose dependence on the distance parameter is (inverse) linear. In contrast, Glasner and Servedio (in *Proceedings of RANDOM*, 2007) prove that every distribution-free testing algorithm for monotone monomials as well as for general monomials must have query complexity $\tilde{\Omega}(n^{1/5})$ (for a constant distance parameter ϵ).

In this paper we present distribution-free testing algorithms for these classes where the query complexity of the algorithms is $\tilde{O}(n^{1/2}/\epsilon)$. We note that as opposed to previous results for distribution-free testing, our algorithms do not build on the algorithms that work under the uniform distribution. Rather, we define and exploit certain structural properties of monomials (and functions that differ from them in a non-negligible manner), which were not used in previous work on property testing.

*Research supported by the Israel Science Foundation (grant No. 246/08)

1 Introduction

Testers (for properties of functions) are algorithms that decide whether a given function has a prespecified property or is “far” from having the property with respect to some fixed distance measure. In most works on property testing, distance is measured with respect to the uniform distribution over the function domain. While in many contexts this distance is appropriate, as it corresponds to assigning equal “importance” or weight to each point in the domain, there are scenarios in which we may want to deal with an underlying weight distribution that is not uniform, and furthermore, is not known to the algorithm. We refer to the latter model as *distribution-free* property testing, while testing under the uniform distribution is considered to be the *standard* model. In the standard model the algorithm is given query access to the tested function and in the distribution-free model the algorithm is also given access to examples distributed according to the unknown underlying distribution.

Indeed, the notion of distribution-free testing is inspired by the distribution-free (Probably Approximately Correct (PAC)) learning model [Val84] and understanding the relation between testing and learning is one of the motivations for property testing. As observed in [GGR98], the complexity of testing a function class \mathcal{F} (that is, testing the property of membership in \mathcal{F}), is not higher than (proper) learning the class \mathcal{F} (under the same conditions, e.g., with respect to the uniform distribution or distribution-free). In view of this, a natural question is for what classes of functions is the complexity of testing *strictly* lower than that of learning. Here, when we refer to the complexity of the algorithm, our main focus is on its query complexity (where in this complexity we count all queries: both on arbitrary points selected by the algorithm, and on points sampled according to the underlying distribution). Note that, as opposed to learning, if we have a testing algorithm for (membership in) a class of functions \mathcal{F} , this does not imply that we have a testing algorithm (with similar complexity) for all subclasses \mathcal{F}' of \mathcal{F} .

There is quite a large variety of function classes for which the complexity of testing is strictly lower than that of learning when the underlying distribution is uniform (e.g., linear functions [BLR93], low-degree polynomials [RS96], singletons, monomials [PRS02] and small monotone DNF [PRS02], monotone functions (e.g., [EKK⁺00, DGL⁺99]), small juntas [FKR⁺04], small decision lists, decision trees and (general) DNF [DLM⁺07] linear threshold functions [MORS09], and more). In contrast, there are relatively few such positive results for distribution-free testing [HK03, HK04, HK07], and, in general, designing distribution-free testing algorithms tends to be more challenging.

One of the main positive results for distribution-free testing [HK07] is that every function class that has a standard tester and can be efficiently *self-corrected* [BLR93], has a distribution-free tester whose complexity is similar to that of the standard tester. In particular this implies that there are efficient distribution-free testers for linear functions and more generally, for low-degree polynomials [HK07]. However, there are function classes of interest (in particular from the point of view of learning theory), which have efficient standard testers, but for which self-correctors do not exist (or are not known to exist). Several such classes (of Boolean functions over $\{0, 1\}^n$) were studied by Glasner and Servedio [GS07]. Specifically, they consider monotone monomials, general monomials, decisions lists, and linear threshold functions. They prove that for these classes, in contrast to standard testing, where the query complexity does not depend on the number of variables n , every distribution-free testing algorithm must make $\Omega((n/\log n)^{1/5})$ queries (for a constant distance parameter ϵ). While these negative results establish that a strong dependence on n is unavoidable for these functions classes in the distribution-free case, it still leaves open the question of whether some *sublinear* dependence on n can be obtained (where distribution-free learning (with queries) requires at least linear complexity [Tur93]).

OUR RESULTS. In this work we prove that both for monotone monomials and for general monomials, a sub-linear dependence on n can be obtained for distribution-free testing. Specifically, we describe distribution-free testing algorithms for these families whose query complexity is $O(\sqrt{n} \log n/\epsilon)$. Thus we advance our knowledge concerning efficient distribution-free testing for two basic function classes. Furthermore, while previous distribution-free testing algorithms are based on, and are similar to the corresponding standard testing algorithms, this is not the case for our algorithms. Rather, we define and exploit certain structural properties of monomials (and functions that differ from them in a non-negligible manner), which were not used in previous work on property testing in the standard model. In what follows we give some intuition concerning the difficulty encountered when trying to extend standard testing of (monotone) monomials to distribution-free testing and then shortly discuss the ideas behind our algorithms.

STANDARD VS. DISTRIBUTION-FREE TESTING OF MONOMIALS. The first simple observation concerning testing monomials under the uniform distribution is the following. If f is a k -monomial (that is, a conjunction of k literals), then $\Pr[f(x) = 1] = 2^{-k}$ (where the probability is over a uniformly selected x). This implies that we can effectively consider only relatively small monomials, that is, k -monomials for which $k = \log(O(1/\epsilon))$, and it allows the testing algorithm to have an exponential dependence on k (since this translates to a linear dependence on $1/\epsilon$). This is not in general the case when the underlying distribution is arbitrary. In particular, the functions considered in the lower bound proof of [GS07] (some of which are monomials, and some of which are far from being monomials), depend on $\Omega(n)$ variables. Thus, for these functions, considering uniformly selected points, essentially gives no information (since the function assigns value 0 to all but a tiny fraction of the points). Furthermore, the support of the distribution D defined in [GS07] is such that the following holds. If one takes a sample (distributed according to D) of size smaller than the square-root of the support size of D , (where there are roughly $n^{2/5}$ points in the support), and performs queries on the sampled points, then it is not possible to distinguish between the monomials and the functions that are far from being monomials (with respect to D). Thus, by sampling according to D , we essentially get no information unless the size of the sample is above a (fairly high) threshold. On the other hand, if we perform queries outside the support of D , then intuitively (and this is formalized in [GS07]), violations (with respect to being a monomial) are hard to find.

Before continuing with a high level description of our algorithms, we note that if we restrict the task of testing to distribution-free testing of (monotone) k -monomials, where k is fixed, then there is an algorithm whose query complexity grows exponentially with k . This follows by combining two results: (1) The aforementioned result of Halevy and Kushilevitz [HK07] concerning the use of “self-correction” in transforming standard testing algorithm to distribution-free testing algorithms; (2) The result of Parnas et al. [PRS02] for testing (monotone) monomials, which has a self-corrector (with complexity 2^k) as a building block. Hence, for small k (i.e., k that is strictly smaller than $\log n$) we have an algorithm with complexity that is sublinear in n . Hence, the question is what can be done when it is not assumed that k is small.

OUR ALGORITHMS: IDEAS AND TECHNIQUES. In what follows we discuss the algorithm for testing monotone monomials over $\{0, 1\}^n$. The algorithm for testing general monomials has the same high-level structure, and can be viewed as a generalization of the algorithm for testing monotone monomials.

We start by introducing the notion of a *violation* hypergraph for a function f . The vertex-set of this hypergraph is $\{0, 1\}^n$, and its edge-set corresponds to subsets that contain evidence that f is not a monotone monomial. Each (hyper)edge includes a single point y^0 such that $f(y^0) = 0$, and for each additional point y^j in the edge, $f(y^j) = 1$. For each such subset there is no monotone monomial that is consistent with f on the subset. For example, we may have $y^0 = 010$, $y^1 = 011$ and $y^2 = 110$ (since y^0 “forces” either x_1

or x_3 to be in the monomial whereas y^1 and y^2 “disallow” these possibilities). Thus, the edge-set of the hypergraph may be exponentially large in n , and edges may have large size (e.g., $\Omega(n)$). Clearly, if f is a monotone monomial, then the hypergraph has no edges. On the other hand, we prove that if f is far from being a monotone monomial (with respect to the underlying distribution, D), then every vertex cover of the (edges of the) hypergraph must have relatively large weight (with respect to D).

Assuming from this point on that f is far from being a monotone monomial (and hence its violation hypergraph has a relatively large weight minimum vertex cover), our algorithm tries to find a small edge in the hypergraph. To this end we do the following. First we take a random sample T of $\Theta(\sqrt{n}/\epsilon)$ points (generated according to D), and consider all points in the sample that are labeled 0 by f . Observe that if f were a monotone monomial, then for each such sample point $y \in f^{-1}(0)$, there must exist at least one index, j , such that x_j is a variable in the monomial and $y_j = 0$. But then $f(1^{j-1}01^{n-j})$ must be 0. In view of this, for each $y \in T \cap f^{-1}(0)$, we search for such an index j (satisfying $y_j = 0$ and $f(1^{j-1}01^{n-j}) = 0$). The search is initiated with the candidate set $\{j : y_j = 0\}$, and the set is cut in half in each iteration (by performing two queries). Thus, if f were a monotone monomial, then such a search must always succeed. On the other hand, if the search fails for any $y \in f^{-1}(0)$, then we obtain evidence that f is not a monotone monomial, and the algorithm may reject. This evidence is an edge (of size 3) in the violation hypergraph (e.g., $(0^t 1^{n-t}, 0^{t/2} 1^{n-t/2}, 1^{t/2} 0^{t/2} 1^{n-t})$ where the first point is in $f^{-1}(0)$ and the latter two are in $f^{-1}(1)$).

Assuming no search fails, the algorithm has a set J of “representative indices”. These indices are such that if f were a monotone monomial, then for each $j \in J$, the variable x_j would be among the variables in f . This means that for every $w \in f^{-1}(1)$ and $j \in J$, it would hold that $w_j = 1$. In the second stage of the algorithm we take an additional sample of $\Theta(\sqrt{n}/\epsilon)$ points and consider all sample points in $f^{-1}(1)$. If for any such sample point w we have that $w_j = 0$ for some $j \in J$, then the algorithm has evidence that f is not a monotone monomial (an edge of size 2 in the violation hypergraph) and it rejects. The crux of the proof is showing that if the probability that the algorithm does not find evidence (in both stages) is small, then it is possible to construct a small-weight vertex cover in the violation hypergraph (implying that f is close to being a monotone monomial).

OTHER RELATED WORK. In addition to the results mentioned previously, Halevy and Kushilevitz [HK03, HK07] study distribution-free testing of monotonicity for functions $f : \Sigma^n \rightarrow R$ (where Σ and R are fully ordered). Building on the (one-dimensional) standard testing algorithm in [EKK⁺00] they give a distribution-free testing algorithm whose query complexity is $O((2 \log |\Sigma|)^n / \epsilon)$. Thus, the dependence on the dimension, n is exponential, in contrast to some of the standard testing algorithms for monotonicity [GGL⁺00, DGL⁺99] where the dependence on n is linear.¹ In follow-up work [HK05, HK07], Halevy and Kushilevitz showed that the exponential dependence on n is unavoidable for distribution-free testing even in the case of Boolean functions over the Boolean hypercube (that is, $|\Sigma| = |R| = 2$).

Halevy and Kushilevitz [HK04] also study distribution-free testing of graph properties in sparse graphs, and give an algorithm for distribution-free testing of connectivity, with similar complexity to the standard testing algorithm for this property.

We note that for some properties that have efficient standard testers, the algorithms can be extended to work under more general families of distributions such as product distributions (e.g., [FKR⁺04, DLM⁺07]). In recent work, Kopparty and Saraf [KS09] consider tolerant testing [PRR06] of linearity under non-uniform distributions (that have certain properties).

¹To be precise, the complexity of the algorithm in [DGL⁺99] is $O(n \log |\Sigma| \log |R| / \epsilon)$, where $|R|$ is the effective size of the range of the function, that is, the number of distinct values of the function.

ORGANIZATION. We start by introducing some notation and definitions in Section 2. In Section 3 we describe and analyze the distribution-free testing algorithm for monotone monomials, and in Section 4 we explain how to extend it to general monomials.

FURTHER RESEARCH. Perhaps the first question that comes to mind is what is the exact complexity of distribution-free testing of (monotone) monomials given the gap between our upper bound and the lower bound of [GS07]. It will also be interesting to design sublinear algorithms for testing the other function classes studied in [GS07]. Another direction is to study testing of monomials and other basic function classes under known distributions (other than the uniform distribution).

2 Preliminaries

For an integer k we let $[k] \stackrel{\text{def}}{=} \{1, \dots, k\}$. In all that follows we consider Boolean functions f whose domain is $\{0, 1\}^n$.

Definition 1 (Monomials) *A function $f : \{0, 1\}^n \rightarrow \{0, 1\}$ is a monomial if it is a conjunction (“and”) of a subset of the literals $\{x_1, \bar{x}_1, \dots, x_n, \bar{x}_n\}$. It is a monotone monomial if it is a conjunction only of variables (and no negations of variables). We denote the class of monomials by \mathcal{MON} and the class of monotone monomials by $\mathcal{MON}_{\mathcal{M}}$.*

We note that we allow the special case that the subset of literals (variables) is empty, in which case f is the all-1 function. In Subsections 3.4 and 4.4 we discuss how to augment our tests so that they work for the case that the subset of literals (variables) must be non-empty.

Definition 2 (Distance) *For two functions $f, g : \{0, 1\}^n \rightarrow \{0, 1\}$ and a distribution D over $\{0, 1\}^n$, we let*

$$\text{dist}_D(f, g) \stackrel{\text{def}}{=} \Pr_{x \sim D}[f(x) \neq g(x)] \quad (1)$$

denote the distance between f and g with respect to D . For a class of Boolean functions \mathcal{F} over $\{0, 1\}^n$ and a function $f : \{0, 1\}^n \rightarrow \{0, 1\}$, we let

$$\text{dist}_D(f, \mathcal{F}) \stackrel{\text{def}}{=} \min_{g \in \mathcal{F}} \{\text{dist}_D(f, g)\} \quad (2)$$

denote the distance between f and the class of functions \mathcal{F} .

Definition 3 (Distribution-Free Testing) *Let \mathcal{F} be a class of Boolean functions over $\{0, 1\}^n$. A distribution-free testing algorithm for (membership in) \mathcal{F} is given access to examples that are distributed according to an unknown distribution D and is given query access to f . The algorithm is also given a distance parameter $0 < \epsilon < 1$, and is required to behave as follows.*

- *If $f \in \mathcal{F}$, then the algorithm should output accept with probability at least $2/3$.*
- *If $\text{dist}_D(f, \mathcal{F}) > \epsilon$, then the algorithm should output reject with probability at least $2/3$.*

If the algorithm accepts every $f \in \mathcal{F}$ with probability 1, then it is a one-sided error algorithm.

In all that follows f always denotes the (unknown) tested function, and D denotes the (unknown) underlying distribution with respect to which the testing algorithm should work. For a point $y \in \{0, 1\}^n$ let $D(y)$ denote the probability assigned to y by D , and for a subset $S \subseteq \{0, 1\}^n$ let $D(S) = \sum_{y \in S} D(y)$ denote the weight that D assigns to the subset S .

We assume without loss of generality that $\epsilon \geq 2^{-n}$, or else, by performing a number of queries that is linear in $1/\epsilon$ (that is, querying f on all domain elements) it is possible to determine whether f is a monotone monomial.

3 Distribution-Free Testing of Monotone Monomials

Recall that in our definition of (monotone) monomials, we allowed the monomial to be “empty” (depend on no variables). That is, the class $\mathcal{MON}_{\mathcal{M}}$ includes the all-1 function. At the end of this subsection we discuss how to augment our test so that it work for the class of all monotone monomials that are a conjunction of at least one variable.

We start by introducing the notion of a *violation hypergraph* of a function and establishing its relation to (the distance to) monotone monomials.

3.1 The Violation Hypergraph

Before defining the violation hypergraph, we introduce some notation. For each point $y \in \{0, 1\}^n$ let $Z(y) \stackrel{\text{def}}{=} \{i : y_i = 0\}$. We use 1^n to denote the all-1 vector (point).

Let g be a Boolean function over $\{0, 1\}^n$ and let $\{y^0, y^1, \dots, y^t\} \subseteq \{0, 1\}^n$ be a subset of points such that $g(y^0) = 0$ and $g(y^j) = 1$ for all $1 \leq j \leq t$. A simple but useful observation is that if g is a monotone monomial, then $Z(y^0)$ must include at least one index i such that $i \notin \bigcup_{j=1}^t Z(y^j)$. This observation motivates the next definition.

Definition 4 (Violation Hypergraph) *Let $H_f = (V(H_f), E(H_f))$ be the hypergraph whose vertex set, $V(H_f)$ is $\{0, 1\}^n$, and whose edge set, $E(H_f)$, contains all subsets $\{y^0, y^1, \dots, y^t\} \subseteq \{0, 1\}^n$ of the following form:*

- $f(y^0) = 0$ and $f(y^j) = 1$ for all $1 \leq j \leq t$.
- $Z(y^0) \subseteq \bigcup_{j=1}^t Z(y^j)$.

For example, if $f(0011) = 0$, $f(0110) = 1$ and $f(1011) = 1$, then $Z(0011) = \{1, 2\}$, $Z(0110) = \{1, 4\}$, and $Z(1011) = \{2\}$, and so $\{0011, 0110, 1011\}$ is an edge in H_f . Note that if $f(1^n) = 0$, then $E(H_f)$ contains the edge $\{y^0 = 1^n\}$ (because $Z(y^0) = \emptyset$ and $\bigcup_{j=1}^t Z(y^j)$ is trivially empty as well).

By the observation preceding Definition 4, if f is a monotone monomial, then $E(H_f) = \emptyset$. We next claim that the reverse implication holds as well, so that we obtain a characterization of monotone monomials that is based on H_f .

Lemma 1 *If $E(H_f) = \emptyset$, then f is a monotone monomial.*

Proof: Suppose that $E(H_f) = \emptyset$, and let $M = \{i : y_i = 1, \forall y \in f^{-1}(1)\}$. Note that since $E(H_f) = \emptyset$, necessarily $f(1^n) = 1$, and so $f^{-1}(1) \neq \emptyset$. Let h be the monotone monomial that is the conjunction of all

x_i such that $i \in M$ (where if M is empty, then h is the all-1 function). We next show that $f(y) = h(y)$ for all $y \in \{0, 1\}^n$.

By the definition of h , for all $y \in f^{-1}(1)$, $h(y) = 1$. To establish that $h(y) = 0$ for all $y \in f^{-1}(0)$, assume in contradiction that there is a point $y \in f^{-1}(0)$ such that $h(y) = 1$. Since $h(y) = 1$ we have (by the definition of h) that $y_i = 1$ for all $i \in M$. It follows that for each i such that $y_i = 0$ there is at least one point $y' \in f^{-1}(1)$ with $y'_i = 0$. That is, $Z(y) \subseteq \bigcup_{y' \in f^{-1}(1)} Z(y')$. But this means that $\{y\} \cup f^{-1}(1)$ is an edge in H_f and we have reached a contradiction. ■

Recall that a vertex cover of a hypergraph is a subset of the vertices that intersects every edge in the hypergraph. We next establish that if f is far from being a monotone monomial (with respect to D), then every vertex cover of H_f must have large weight (with respect to D). This lemma strengthens Lemma 1 in the following sense. Lemma 1 is equivalent to saying that if f is not a monotone monomial, then $E(H_f) \neq \emptyset$. In particular this implies that if f is not a monotone monomial, then every vertex cover of H_f is non-empty. Lemma 2 can be viewed as quantifying this statement (and taking into account the underlying distribution D).

Lemma 2 *If $\text{dist}_D(f, \mathcal{MON}_{\mathcal{M}}) > \epsilon$, then for every vertex cover C of H_f we have $D(C) > \epsilon$.*

Proof: The proof of Lemma 2 follows by observing that Lemma 1 and its proof can be easily extended to any sub-hypergraph of H_f that is determined by some subset $R \subseteq \{0, 1\}^n$. For the sake of completeness, we next give the details.

Assume, contrary to the claim, that there exists a vertex cover C of H_f such that $D(C) \leq \epsilon$. Let $R = \{0, 1\}^n \setminus C$ be the vertices that do not belong to the vertex-cover. Let $H_f(R) = (R, E(H_f(R)))$ be the sub-hypergraph of H_f where $E(H_f(R))$ consists of all edges in H_f that are contained² in R . Since C is a vertex cover, we have that $E(H_f(R)) = \emptyset$. We next define a monotone monomial h and show that it is consistent with f on R . Since $D(C) \leq \epsilon$ this implies that $\text{dist}_D(h, f) \leq \epsilon$, in contradiction to the premise of the lemma.

If $f^{-1}(1) \cap R = \emptyset$, then, since $E(H_f(R)) = \emptyset$, necessarily $1^n \notin R$. In this case we let h be the monomial that is the conjunction of all variables x_i (so that it has value 1 only on 1^n). Since $f^{-1}(1) \cap R = \emptyset$, this monomial is consistent with all points in R . Otherwise, let $M = \{i : y_i = 1, \forall y \in f^{-1}(1) \cap R\}$. Let h be the monotone monomial that is the conjunction of all x_i such that $i \in M$ (where if M is empty, then h is the all-1 function). We next show that $f(y) = h(y)$ for all $y \in \{0, 1\}^n \cap R$.

By the definition of h , for all $y \in f^{-1}(1) \cap R$, $h(y) = 1$. To establish that $h(y) = 0$ for all $y \in f^{-1}(0) \cap R$, assume in contradiction that there is a point $y \in f^{-1}(0) \cap R$ such that $h(y) = 1$. Since $h(y) = 1$ we have (by the definition of h) that $y_i = 1$ for all $i \in M$. It follows that for each i such that $y_i = 0$ there is at least one point $y' \in f^{-1}(1) \cap R$ with $y'_i = 0$. That is, $Z(y) \subseteq \bigcup_{y' \in f^{-1}(1) \cap R} Z(y')$. But this means that $\{y\} \cup (f^{-1}(1) \cap R)$ is an edge in $H_f(R)$ and we have reached a contradiction. ■

By Lemmas 1 and 2, if f is a monotone monomial, then $E(H_f) = \emptyset$, so that trivially every minimum vertex cover of H_f is empty, while if $\text{dist}_D(f, \mathcal{MON}_{\mathcal{M}}) > \epsilon$, then every vertex cover of H_f has weight greater than ϵ with respect to D . We would like to show that this implies that if $\text{dist}_D(f, \mathcal{MON}_{\mathcal{M}}) > \epsilon$, then we can actually find (with high probability) an edge in H_f , which provides evidence to the fact that f is not a monotone monomial.

²We could refer to this as the sub-hypergraph *induced* by R , but this term is used for a slightly different notion in the context of hypergraphs.

3.2 The Testing Algorithm

We first introduce a few more notation. Let $\bar{e}^i = 1^{i-1}01^{n-i}$. For any subset $Z \subseteq [n]$, let $y(Z)$ be the point in $\{0, 1\}^n$ such that for every $i \in Z$ its i^{th} coordinate is 1, and for every $i \notin Z$ its i^{th} coordinate is 0. For any subset $S \subseteq \{0, 1\}^n$, let $S_{f,0} = \{y \in S : f(y) = 0\}$ and $S_{f,1} = \{y \in S : f(y) = 1\}$.

The first observation on which our algorithm is based is that for every point $y \in f^{-1}(0)$, there must be at least one index $i \in Z(y)$ for which $f(\bar{e}^i) = 0$, or else we have evidence that f is not a monotone monomial. In fact, we don't need to verify that $f(\bar{e}^i) \neq 0$ for every $i \in Z(y)$ in order to obtain evidence that f is not a monotone monomial. Rather, if we search for such an index (in a manner described momentarily), and this search fails, then we already have evidence that f is not a monotone monomial.

The search procedure (which performs a binary search), receives as input a point $y \in f^{-1}(0)$ and searches for an index $j \in Z(y)$ such that $f(\bar{e}_j) = 0$. This is done by repeatedly partitioning a set of indices, Z , starting with $Z = Z(y)$, into two parts Z_1 and Z_2 of (almost) equal size, and continuing the search with a part Z_i , $i \in \{1, 2\}$ for which $f(y(Z_i)) = 0$. (If both parts satisfy the condition, then we continue with Z_1 .) Note that if both $f(y(Z_1)) = 1$ and $f(y(Z_2)) = 1$, then we have evidence that f is not a monotone monomial because $f(y(Z_1 \cup Z_2)) = 0$ (so that $\{y(Z_1 \cup Z_2), y(Z_1), y(Z_2)\}$ is an edge in H_f). The search also fails (from the start) if $Z(y) = \emptyset$ (that is, $y = 1^n$). For the precise pseudo-code of the procedure, see Figure 1.

Algorithm 1: Binary Search (Input: $y \in \{0, 1\}^n$)

1. $Z \leftarrow Z(y)$.
2. if $|Z| = 0$, then output fail and halt.
3. While ($|Z| \geq 2$) do
 - (a) Let (Z_1, Z_2) be a fixed partition of Z where $||Z_1| - |Z_2|| \leq 1$.
Specifically, Z_1 is the set of the first $\lfloor |Z|/2 \rfloor$ indices in Z .
 - If $f(y(Z_1)) = 0$, then $Z \leftarrow Z_1$;
 - else if $f(y(Z_2)) = 0$, then $Z \leftarrow Z_2$;
 - else output fail and halt.
4. Output the single index that remains in Z .

Figure 1: The binary search procedure for monotone monomials.

The testing algorithm starts by obtaining a sample of $\Theta(\sqrt{n}/\epsilon)$ points, where each point is generated independently according to D . (Since the points are generated independently, repetitions may occur.) For each point in the sample that belongs to $f^{-1}(0)$, the algorithm calls the binary search procedure. If any search fails, then the algorithm rejects f (recall that in such a case the algorithm has evidence that f is not a monotone monomial). Otherwise, the algorithm has a collection of indices J such that $f(\bar{e}_j) = 0$ for every $j \in J$. The algorithm then takes an additional sample, also of size $\Theta(\sqrt{n}/\epsilon)$, and checks whether there exists a point y in the sample such that $f(y) = 1$ and $Z(y)$ contains some $j \in J$. In such a case the algorithm has evidence that f is not a monotone monomial (specifically, $\{\bar{e}_j, y\}$ is an edge in H_f), and it rejects. For the precise pseudo-code of the algorithm, see Figure 2.

We shall use Lemma 2 to show that if $\text{dist}_D(f, \text{MON}_{\mathcal{M}})$ is relatively large, then either the first sample

will contain a point on which the binary search procedure fails (with high probability over the choice of the first sample), or the second sample will contain a point y such that $f(y) = 1$ and $Z(y) \cap J \neq \emptyset$ (with high probability over the choice of both samples).

Algorithm 2: Monotone Monomials Test

1. Obtain a sample T of $\Theta(\sqrt{n}/\epsilon)$ points, each generated independently according to D .
2. For each point $y \in T_{f,0}$ run the binary search procedure (Algorithm 1) on y .
3. If the binary search fails for any of the points, then output reject and halt. Otherwise, for each $y \in T_{f,0}$ let $j(y)$ be the index returned for y , and let $J(T_{f,0}) = \bigcup_{y \in T_{f,0}} j(y)$.
4. Obtain another sample T' of size $\Theta(\sqrt{n}/\epsilon)$ (generated independently according to D).
5. If there is a point $y \in T'_{f,1}$ such that $Z(y) \cap J(T_{f,0}) \neq \emptyset$, then output reject, otherwise output accept.

Figure 2: The distribution-free testing algorithm for monotone monomials.

3.3 The Analysis of the Testing Algorithm for Monotone Monomials

The next definition will serve us in the analysis of the algorithm.

Definition 5 (Empty points and representative indices) For a point $y \in f^{-1}(0)$, we say that y is empty (with respect to f) if the binary search procedure (Algorithm 1) fails on y . We denote the set of empty points (with respect to f) by $Y_\emptyset(f)$. If y is not empty, then we let $j(y) \in Z(y)$ denote the index that the binary search procedure returns. We refer to this index as the representative index for y . If $y \in Y_\emptyset(f)$, then $j(y)$ is defined to be 0.

Note that since the binary search procedure is deterministic, the index $j(y)$ is uniquely defined for each $y \notin Y_\emptyset(f)$.

As in Algorithm 2, for a sample T and $T_{f,0} = T \cap f^{-1}(0)$, we let $J(T_{f,0}) = \{j(y) : y \in T_{f,0} \setminus Y_\emptyset(f)\}$ denote the set of representative indices for the sample. For any subset $J \subseteq [n]$, let $Y_{f,1}(J)$ denote the set of all points $y \in f^{-1}(1)$ for which $Z(y) \cap J \neq \emptyset$. In particular, if we set $J = J(T_{f,0})$, then each point $y \in Y_{f,1}(J)$, together with any index j in its intersection with J , provide evidence that f is not a monotone monomial (i.e., $\{\bar{e}_j, y\} \in E(H_f)$). We next state our main lemma.

Lemma 3 Suppose that $\text{dist}_D(f, \text{MON}_{\mathcal{M}}) > \epsilon$ and consider a sample T of $c_1\sqrt{n}/\epsilon$ points generated independently according to D . For a sufficiently large constant c_1 , with probability at least $5/6$ over the choice of T , either $T_{f,0}$ contains an empty point (with respect to f) or $D(Y_{f,1}(J(T_{f,0}))) \geq \frac{\epsilon}{4\sqrt{n}}$.

The proof of Lemma 3 builds on Claim 4, stated next, which in turn uses the following notation: For a subset $J \subseteq [n]$, we let $Y_{f,0}(J)$ denote the set of points $y \in f^{-1}(0)$ for which³ $j(y) \in J$.

³Note that $Y_{f,0}(J)$ and $Y_{f,1}(J)$ do not only differ in that they refer to points in $f^{-1}(0)$ and $f^{-1}(1)$, respectively. Rather, $Y_{f,0}(J)$ is the subset of points y (in $f^{-1}(0)$) whose representative index $j(y)$ belongs to J , while $Y_{f,1}(J)$ is the subset of points y (in $f^{-1}(1)$) for which some index $j \in J$ belongs to J .

Claim 4 Let I be any fixed subset of $[n]$, and consider a sample T of $s = c_1\sqrt{n}/\epsilon$ points generated independently according to D . For a sufficiently large constant c_1 , with probability at least $9/10$ over the choice of T , either $|J(T_{f,0}) \setminus I| \geq \sqrt{n}$ or $D(Y_{f,0}([n] \setminus (J(T_{f,0}) \cup I))) \leq \epsilon/2$.

To make Claim 4 more concrete, consider the special case in which $I = \emptyset$. The lemma simply says that with probability at least $9/10$ (over the choice of T) either the subset of indices $J(T_{f,0})$ is relatively large, or the weight of the set of points y in $f^{-1}(0)$ for which $j(y)$ is not contained in $J(T_{f,0})$ is relatively small.

Proof: Consider selecting the points in T one after the other, where for $\ell = 1, \dots, s = c_1\sqrt{n}/\epsilon$ we let y^ℓ denote the ℓ^{th} point selected, and we let $T^\ell = \{y^1, \dots, y^\ell\}$ (where $T^0 = \emptyset$). Let $Y^\ell = Y_{f,0}([n] \setminus (J(T_{f,0}^{\ell-1}) \cup I))$ denote the set of points in $f^{-1}(0)$ whose representative index differs from all the previously found representative indices and does not belong to I . For each $1 \leq \ell \leq s$, let χ^ℓ be a 0/1 valued random variable, where $\chi^\ell = 1$ if either $y^\ell \in Y^\ell$ (so that $j(y^\ell)$ is a *new* representative index), or $D(Y^\ell) \leq \epsilon/2$. Observe that if for some ℓ_0 we have that $D(Y^{\ell_0}) \leq \epsilon/2$, then $D(Y^\ell) \leq \epsilon/2$ for all $\ell \geq \ell_0$.

While χ^1, \dots, χ^s are not independent random variables, the probability that their sum is smaller than \sqrt{n} (which is less than half the expected value for $c_1 \geq 4$), can be bounded using a multiplicative Chernoff bound that is applied to a similar sequence of *independent* random variables. The basic observation is that the probability that $\chi^\ell = 1$ is greater than $\epsilon/2$ *no matter what were the choices of* $y^1, \dots, y^{\ell-1}$ (which determine $\chi^1, \dots, \chi^{\ell-1}$). Therefore, let $\tilde{\chi}^1, \dots, \tilde{\chi}^s$ be *independent* 0/1 valued variables, where $\Pr[\tilde{\chi}^\ell = 1] = \epsilon/2$ for each $1 \leq \ell \leq s$. Letting $c_1 = 64$ (so that $s = 64\sqrt{n}/\epsilon$), we have that

$$\begin{aligned} \Pr \left[\sum_{\ell=1}^s \chi^\ell < \sqrt{n} \right] &\leq \Pr \left[\sum_{\ell=1}^s \tilde{\chi}^\ell < \sqrt{n} \right] \\ &< \Pr \left[\sum_{\ell=1}^s \tilde{\chi}^\ell < (1/2)(\epsilon/2)s \right] \\ &< \exp(-\epsilon s/16) < 1/6. \end{aligned} \tag{3}$$

This means that with probability at least $9/10$, either $D(Y^{\ell-1}) \leq \epsilon/2$ for some ℓ , in which case $D(Y_{f,0}([n] \setminus (J(T_{f,0}) \cup I))) \leq \epsilon/2$, or $|J(T_{f,0}) \setminus I| \geq \sqrt{n}$. ■

Proof of Lemma 3: Assume, contrary to the claim, that with probability at least $1/6$ over the choice of T (which consist of $c_1\sqrt{n}/\epsilon$ points selected independently according to D), there is no point in T that is empty with respect to f and $D(Y_{f,1}(J(T_{f,0}))) < \frac{\epsilon}{4\sqrt{n}}$. We will show how, using this assumption, it is possible to prove that there exists a vertex cover C of H_f with $D(C) \leq \epsilon$. By Claim 2, this contradicts the fact that $\text{dist}_D(f, \mathcal{MON}_M) > \epsilon$.

THE CONSTRUCTION OF C . We show how to construct a vertex cover C of H_f in three stages. In the first stage we put in C all empty points, that is, all points in $Y_\emptyset(f)$. By the counter-assumption, $D(Y_\emptyset(f)) \leq 2\epsilon/(c_1\sqrt{n})$. This is true because otherwise, the probability that the sample T does not contain an empty point is at most $(1 - 2\epsilon/(c_1\sqrt{n}))^{c_1\sqrt{n}/\epsilon} < e^{-2} < 1/6$.

In the second stage we work in iterations. In each iteration ℓ we add to the cover a subset of points $Y^\ell \subseteq f^{-1}(1)$, which is determined by a subset $T^\ell \subseteq \{0, 1\}^n$. These subsets are determined as follows. Let $I^1 = \emptyset$, and for $\ell > 1$ let $I^\ell = I^{\ell-1} \cup J(T_{f,0}^{\ell-1})$. We shall think of I^ℓ as the subset of representative indices that have “already been covered” in a sense that will become clear momentarily. Suppose we apply Claim 4 with $I = I^\ell$. The claim says that with probability at least $9/10$ over the choice of T , either $|J(T_{f,0}) \setminus I^\ell| \geq \sqrt{n}$ (that is, there are many “new” representative indices corresponding to points in $T_{f,0}$),

or $D(Y_{f,0}([n] \setminus (J(T_{f,0}) \cup I^\ell))) \leq \epsilon/2$ (that is, the total weight of the points whose representative index is not already in I^ℓ or $J(T_{f,0})$ is small). Thus, the probability that neither of the two hold is at most $1/10$.

Combining this with our counter-assumption (and taking a union bound), we have that with probability at least $1/6 - 1/10 > 0$ over the choice of T : $D(Y_{f,1}(J(T_{f,0}))) < \frac{\epsilon}{4\sqrt{n}}$ (that is, the total weight of the points y in $f^{-1}(1)$ such that $Z(y) \cap J(T_{f,0}) \neq \emptyset$ is small), and either $|J(T_{f,0}) \setminus I^\ell| \geq \sqrt{n}$ or $D(Y_{f,0}([n] \setminus (J(T_{f,0}) \cup I^\ell))) \leq \epsilon/2$. Since this (combined) event occurs with probability greater than 0, there must exist at least one set T for which it holds. Denote this set by T^ℓ , and let $Y^\ell = Y_{f,1}(J(T_{f,0}^\ell))$ (so that all points in Y^ℓ are added to the cover). If $D(Y_{f,0}([n] \setminus (J(T_{f,0}^\ell) \cup I^\ell))) \leq \epsilon/2$, then this (second) stage ends, and in the third stage we add to the cover C all points in $Y_{f,0}([n] \setminus (J(T_{f,0}^\ell) \cup I^\ell))$. Otherwise, we set $I^{\ell+1} = I^\ell \cup J(T_{f,0}^\ell)$ and continue with the next iteration.

ESTABLISHING THAT C IS A VERTEX COVER OF H_f . Consider any point $y \in f^{-1}(0)$ that is contained in at least one edge in H_f . We shall show that either $y \in C$, or, for every edge $B \in H_f$ that contains y , there is at least one point $y' \in B \cap f^{-1}(1)$ that belongs to C . Since each edge in H_f contains some point $y \in f^{-1}(0)$, this implies that C is a vertex cover. Details follow.

If $y \in Y_\emptyset(f)$, then it is added to the cover in the first stage. Otherwise, y has a representative index $j(y) \in Z(y)$. Consider the iterations in the second stage of the construction of C . If for some iteration ℓ we have that $j(y) \in J(T_{f,0}^\ell)$ (where we consider the first iteration in which this occurs), then the cover contains all points in $Y_{f,1}(\{j(y)\}) \subseteq Y_{f,1}(J(T_{f,0}^\ell)) = Y^\ell$. But since, by the definition of H_f , each edge in H_f that contains y must contain some $y' \in f^{-1}(1)$ such that $j(y) \in Z(y')$, all edges containing y are covered after iteration ℓ . On the other hand, if $j(y) \notin J(T_{f,0}^\ell)$ for every iteration ℓ , then let ℓ^* denote the index of the last iteration and consider the third stage. In this stage we add to C all points in $Y_{f,0}([n] \setminus (J(T_{f,0}^{\ell^*}) \cup I^{\ell^*}))$. But this set consists of those points whose representative index is not contained in $J(T_{f,0}^{\ell^*}) \cup I^{\ell^*} = \bigcup_{\ell=1}^{\ell^*} J(T_{f,0}^\ell)$, and in particular the set contains y , so that y is added to the cover in the third stage.

BOUNDING THE WEIGHT OF C . The main observation is that in each iteration ℓ of the second stage except, possibly, for the last one (ℓ^*), we have that $|J(T_{f,0}^\ell) \setminus I^\ell| \geq \sqrt{n}$. That is, each such iteration corresponds to a subset of at least \sqrt{n} indices in $[n]$, where the different subsets are disjoint. This implies that $\ell^* \leq \sqrt{n} + 1$. By the construction of C ,

$$\begin{aligned} D(C) &= D(Y_\emptyset(f)) + \sum_{\ell=1}^{\ell^*} D\left(Y_{f,1}(J(T_{f,0}^\ell))\right) + D\left(Y_{f,0}([n] \setminus (J(T_{f,0}^{\ell^*}) \cup I^{\ell^*}))\right) \\ &\leq \frac{2\epsilon}{c_1\sqrt{n}} + (\sqrt{n} + 1) \cdot \frac{\epsilon}{4\sqrt{n}} + \epsilon/2 \leq \epsilon, \end{aligned} \quad (4)$$

where the last inequality holds for $c_1 \geq 8$ (assuming $n \geq 4$). ■

Theorem 5 *Algorithm 2 is a distribution-free 1-sided-error testing algorithm for (membership in) $\mathcal{MON}_{\mathcal{M}}$. Its query complexity is $O(\sqrt{n} \log n/\epsilon)$.*

Proof: Consider first the case that f is a monotone monomial. Observe that the algorithm rejects only if it finds evidence that f is not a monotone monomial. This evidence is either in the form of two (disjoint) subsets of indices, Z_1 and Z_2 such that $f(y(Z_1)) = f(y(Z_2)) = 1$ while $f(y(Z_1 \cup Z_2)) = 0$ (found by the binary search procedure), or it is of the form of an index j and a point $y \in f^{-1}(1)$, such that $f(\bar{e}_j) = 0$ and $j \in Z(y)$. Therefore, the algorithm never rejects a monotone monomial.

Consider next the case that $\text{dist}_D(f, \mathcal{MON}_{\mathcal{M}}) > \epsilon$. By Lemma 3, for a sufficiently large constant c_1 in the $\Theta(\cdot)$ notation for T (the first sample), with probability at least $5/6$ over the choice of T , either there is an empty point in $T_{f,0} \subseteq T$, or $D(Y_{f,1}(J(T_{f,0}))) \geq \frac{\epsilon}{4\sqrt{n}}$. If there is an empty point in $T_{f,0}$, then the binary search will fail on that point and the algorithm will reject. On the other hand, if $D(Y_{f,1}(J(T_{f,0}))) \geq \frac{\epsilon}{4\sqrt{n}}$, then, since the size of the second sample, T' , is $c'_1\sqrt{n}/\epsilon$, the probability that no point $y \in Y_{f,1}(J(T_{f,0}))$ is selected in T' is at most $(1 - \frac{\epsilon}{4\sqrt{n}})^{c'_1\sqrt{n}/\epsilon}$, which is upper bounded by $1/6$ for $c'_1 \geq 8$. But if such a point is selected, then the algorithm rejects.⁴ Therefore, the probability that the algorithm rejects a function f for which $\text{dist}_D(f, \mathcal{MON}_{\mathcal{M}}) > \epsilon$ is at least $2/3$.

Finally, the number of points sampled is $O(\sqrt{n}/\epsilon)$ since the algorithm obtains two samples of this size. Since for each point in the first sample that belongs to $f^{-1}(0)$ the algorithm performs a binary search, the query complexity of the algorithm is $O(\sqrt{n} \log n/\epsilon)$. ■

3.4 Disallowing the all-1 function

Our algorithm as described in Subsection 3.2 works for the class of monotone monomials, $\mathcal{MON}_{\mathcal{M}}$, when it is assumed to contain the “empty” monomial, that is, the all-1 function. Let $\mathbf{1}$ denote the all-1 function (over $\{0, 1\}^n$) and let $\mathcal{MON}'_{\mathcal{M}}$ denote the class of monotone monomials that are a conjunction of at least one variable. Thus, $\mathcal{MON}'_{\mathcal{M}} = \mathcal{MON}_{\mathcal{M}} \setminus \{\mathbf{1}\}$. We next show how to augment our testing algorithm for $\mathcal{MON}_{\mathcal{M}}$ (Algorithm 2) so that it work for $\mathcal{MON}'_{\mathcal{M}}$.

First we run Algorithm 2 with the distance parameter ϵ set to $\epsilon/2$ and with slightly larger constants in the sizes of the sample, so that its failure probability is at most $1/6$ rather than $1/3$. If the algorithm rejects, then we reject as well. Otherwise, it is possible that the algorithm accepted f because $\text{dist}_D(f, \mathbf{1}) \leq \epsilon/2$ (so that $\text{dist}_D(f, \mathcal{MON}_{\mathcal{M}}) \leq \epsilon/2$) while $\text{dist}_D(f, \mathcal{MON}'_{\mathcal{M}}) > \epsilon$. We are interested in detecting this (with high probability) and rejecting f in such a case.

To this end we take an additional sample R of $\Theta(\log n/\epsilon)$ points, each generated independently according to D . We then check whether there exists at least one index $i \in [n]$ such that $y_i = 1$ for all points $y \in S \cap f^{-1}(1)$. If there is no such index, then we reject, otherwise we accept. We next show that if $\text{dist}_D(f, \mathbf{1}) \leq \epsilon/2$ and $\text{dist}_D(f, \mathcal{MON}'_{\mathcal{M}}) > \epsilon$, then this simple procedure rejects f with high constant probability.

The simple observation is that if $\text{dist}_D(f, \mathbf{1}) \leq \epsilon/2$ and $\text{dist}_D(f, \mathcal{MON}'_{\mathcal{M}}) > \epsilon$, then $\Pr_{y \sim D}[f(y) = 1 \text{ and } y_i = 0] \geq \epsilon/2$ for every $i \in [n]$. Assume, contrary to the claim, that there exists some index $i \in [n]$ for which $\Pr_{y \sim D}[f(y) = 1 \text{ and } y_i = 0] < \epsilon/2$. Since $\text{dist}_D(f, \mathbf{1}) \leq \epsilon/2$ (so that $\Pr_{y \sim D}[f(y) = 0] \leq \epsilon/2$), we have that $\Pr_{y \sim D}[f(y) = 0 \text{ and } y_i = 1] \leq \epsilon/2$. Therefore, by flipping the value of f on all points y such that either $f(y) = 1$ and $y_i = 0$ or $f(y) = 0$ and $y_i = 1$ we obtain a monotone monomial (singleton) g such that $\text{dist}_D(f, g) \leq \epsilon$, and we have reached a contradiction.

Finally, if $\Pr_{y \sim D}[f(y) = 1 \text{ and } y_i = 0] \geq \epsilon/2$ for every $i \in [n]$, then by taking a union bound over all $i \in [n]$ we get the following. With high constant probability over the choice of a sample of size $\Theta(\log n/\epsilon)$, for each $i \in [n]$, the sample will contain a point y such that $f(y) = 1$ and $y_i = 0$.

⁴We note that the analysis doesn't explicitly address the case that $T_{f,0} = \emptyset$, where the algorithm accepts (for every T') simply because $J(T_{f,0})$ is empty. What the analysis implies (implicitly) is that the probability that such an event occurs when $\text{dist}_D(f, \mathcal{MON}_{\mathcal{M}}) > \epsilon$ is at most a small constant. It is possible to argue this directly (since if $\text{dist}_D(f, \mathcal{MON}_{\mathcal{M}}) > \epsilon$, then in particular, f is ϵ -far with respect to D from the all-1 function, so that the probability that no point in $f^{-1}(0)$ is selected in the first sample is very small).

4 Distribution-Free Testing of (General) Monomials

The high-level structure of the algorithm for testing general monomials is similar to the algorithm for testing monotone monomials, but several modifications have to be made (and hence the algorithm and the notions it is based on are seemingly more complex). In this section we explain what the modifications are and how this affects the analysis.

Recall that for a point $y \in \{0, 1\}^n$ we let $Z(y) = \{i : y_i = 0\}$. Analogously, we let $O(y) = \{i : y_i = 1\}$. For a non-empty set of points $Y \subseteq \{0, 1\}^n$, let $Z(Y) = \bigcap_{y \in Y} Z(y)$ and $O(Y) = \bigcap_{y \in Y} O(y)$. We shall use the convention that $Z(\emptyset) = O(\emptyset) = [n]$.

4.1 The Violation Hypergraph (for General Monomials)

A basic observation concerning (general) monomials is that if g is a monomial, then for every subset $\{y^0, y^1, \dots, y^t\}$ such that $g(y^0) = 0$ and $g(y^j) = 1$ for each $1 \leq j \leq t$, the following holds: There must be at least one index $i \in [n]$ such that either $i \in Z(\{y^1, \dots, y^t\})$ and $y_i^0 = 1$ or $i \in O(\{y^1, \dots, y^t\})$ and $y_i^0 = 0$. This implies that if we have a subset $\{y^0, y^1, \dots, y^t\}$ such that $g(y^0) = 0$ and $g(y^j) = 1$ for each $1 \leq j \leq t$, and such that $Z(\{y^1, \dots, y^t\}) \subseteq Z(y^0)$ and $O(\{y^1, \dots, y^t\}) \subseteq O(y^0)$, then we have evidence that g is not a monomial. This motivates the next (modified) definition of a violation hypergraph.

Definition 6 (Violation hypergraph (w.r.t. general monomials)) *Let $H_f = (V(H_f), E(H_f))$ be the hypergraph whose vertex set, $V(H_f)$ is $\{0, 1\}^n$, and whose edge set, $E(H_f)$, contains all subsets $\{y^0, y^1, \dots, y^t\} \subseteq \{0, 1\}^n$ of the following form:*

- $f(y^0) = 0$ and $f(y^j) = 1$ for all $1 \leq j \leq t$.
- $Z(\{y^1, \dots, y^t\}) \subseteq Z(y^0)$ and $O(\{y^1, \dots, y^t\}) \subseteq O(y^0)$.

Observe that the second item in Definition 4 (of the violation hypergraph for monotone monomials), which requires that $Z(y^0) \subseteq \bigcup_{j=1}^t Z(y^j)$, is equivalent to $O(\{y^1, \dots, y^t\}) \subseteq O(y^0)$. Therefore, the difference between Definition 4 and Definition 6 is in the additional requirement that $Z(\{y^1, \dots, y^t\}) \subseteq Z(y^0)$.

Similarly to Lemma 1 here we have the next lemma.

Lemma 6 *If $E(H_f) = \emptyset$ and $f^{-1}(1) \neq \emptyset$, then f is a monomial.*

Note that slightly differently from Lemma 1, in Lemma 6 we explicitly added the condition that $f^{-1}(1) \neq \emptyset$. The reason is that while in the case of monotone monomials, the fact that $E(H_f) = \emptyset$ implies that $f^{-1}(1) \neq \emptyset$ (because if $f(1^n) = 0$, then $E(H_f)$ is not empty since it contains 1^n), in the case of general monomials this implication does not hold. The proof of Lemma 6 is very similar to the proof of Lemma 1. Essentially the only difference is that we define h to be the conjunction of all variables x_i such that $i \in Z(f^{-1}(1))$ and all negations of variables \bar{x}_i such that $i \in O(f^{-1}(1))$. (In case both subsets are empty we get the all-1 function.) Here too h agrees with f , by its definition, on all points in $f^{-1}(1)$ and it remains to verify that h also agrees with f on all points in $f^{-1}(0)$, using the premise of the lemma that $E(H_f) = \emptyset$.

The next lemma is analogous to Lemma 2 (and is proved very similarly).

Lemma 7 *If $\text{dist}_D(f, \mathcal{MON}) > \epsilon$ and $f^{-1}(1) \neq \emptyset$, then for every vertex cover C of H_f we have $D(C) > \epsilon$.*

4.2 The Algorithm for Testing General Monomials

For a vector $y \in \{0, 1\}^n$, and for an index $i \in [n]$, let y^{-i} be the same as y except that the i^{th} coordinate in y is flipped. That is, $y_\ell^{-i} = y_\ell$ for all $\ell \neq i$ and $y_i^{-i} = \bar{y}_i$. For a subset $I \subseteq [n]$ let y^{-I} be the vector y with each coordinate $i \in I$ flipped. That is, $y_\ell^{-I} = y_\ell$ for all $\ell \notin I$ and $y_\ell^{-I} = \bar{y}_\ell$ for all $\ell \in I$. Let $\Delta(y, w) \subseteq [n]$ be the subset of indices i such that $y_i \neq w_i$, and note that $y = w^{-\Delta}$ for $\Delta = \Delta(y, w)$.

Algorithm 3: Binary Search for General Monomials (Input: $y \in f^{-1}(1), w \in f^{-1}(0)$)

1. $\Delta \leftarrow \Delta(y, w)$;
2. *While* ($|\Delta| \geq 2$) *do*
 - (a) *Let* (Δ_1, Δ_2) *be a fixed partition of* Δ *where* $\|\Delta_1\| - \|\Delta_2\| \leq 1$.
Specifically, Δ_1 is the set of the first $\lfloor |\Delta|/2 \rfloor$ indices in Δ .
 - (b) *If* $f(w^{-\Delta_1}) = 0$, *then* $\Delta \leftarrow \Delta_1$;
 - (c) *else if* $f(w^{-\Delta_2}) = 0$, *then* $\Delta \leftarrow \Delta_2$;
 - (d) *else output fail and halt.*
3. *Output the single index* $j \in \Delta$;

Figure 3: The binary search procedure for general monomials.

We start by describing the binary search procedure (for general monomials). Its pseudo-code is given in Algorithm 3 (see Figure 3). The procedure receives as input two points $w, y \in \{0, 1\}^n$ such that $f(w) = 1$ and $f(y) = 0$ and outputs an index $j \in [n]$ such that $y_j \neq w_j$ and such that $f(w^{-j}) = 0$. If f is a monomial, then at least one such index must exist. Note that if $w = 1^n$, then the output of the search is as specified by the binary search procedure for monotone monomials (Algorithm 1). In fact, Algorithm 1 itself (and not only its output specification) is essentially the same as Algorithm 3 for the special case of $w = 1^n$. (Since $f(1^n)$ must equal 1 if f is a monotone monomial, we can think of the binary search procedure for monotone monomials as implicitly working under this assumption.)

The search is performed by repeatedly partitioning a set of indices Δ , starting with $\Delta = \Delta(y, w)$, into two parts Δ_1 and Δ_2 of (almost) equal size, and querying f on the two points, $w^{-\Delta_1}$ and $w^{-\Delta_2}$. If f returns 1 for both, then the search fails. Otherwise, the search continues with Δ_i for which $f(w^{-\Delta_i}) = 0$, unless $|\Delta_i| = 1$, in which case the desired index is found. If the search fails, then we have evidence that f is not a monomial. Namely, we have three points, $w^{-\Delta_1}$, $w^{-\Delta_2}$ and $w^{-\Delta}$, where $\Delta = \Delta_1 \cup \Delta_2$, such that $f(w^{-\Delta}) = 0$ and $f(w^{-\Delta_1}) = f(w^{-\Delta_2}) = 1$. Since $w^{-\Delta_1}$ and $w^{-\Delta_2}$ disagree on all coordinates in Δ , and all three points agree on all coordinates in $[n] \setminus \Delta$, we have that $Z(\{w^{-\Delta_1}, w^{-\Delta_2}\}) \subseteq Z(w^{-\Delta})$ and $O(\{w^{-\Delta_1}, w^{-\Delta_2}\}) \subseteq O(w^{-\Delta})$, so that the three points constitute an edge in H_f .

The testing algorithm for general monomials starts by obtaining a sample of $\Theta(1/\epsilon)$ points, each generated independently according to D . The algorithm arbitrary selects a point w in this sample that belongs to $f^{-1}(1)$. If no such point exists, then the algorithm simply accepts f (and halts). Otherwise, this point serves as a kind of *reference point*. As in the case of the binary search procedure, the testing algorithm for monotone monomials (Algorithm 2) is essentially the same as the testing algorithm for general monomials (Algorithm 4) with w (implicitly) set to be 1^n .

Next, the algorithm obtains a sample of $\Theta(\sqrt{n}/\epsilon)$ points (each generated independently according to

Algorithm 4: General Monomials Test

1. Obtain a sample S of $\Theta(1/\epsilon)$ points, each generated independently according to D .
2. If $S_{f,1} = \emptyset$, then output accept and halt. Otherwise, arbitrarily select a point $w \in S_{f,1}$.
3. Obtain a sample T of $\Theta(\sqrt{n}/\epsilon)$ points, each generated independently according to D .
4. For each point $y \in T_{f,0}$ run the binary search procedure (Algorithm 3) on w, y .
5. If the binary search fails for any of the points, then output reject and halt. Otherwise, for each $y \in T_{f,0}$ let $j^w(y)$ be the index returned for y , and let $J^w(T_{f,0}) = \{j^w(y) : y \in T_{f,0}\}$.
6. Obtain another sample T' of size $\Theta(\sqrt{n}/\epsilon)$ (generated independently according to D).
7. If there is a point $y \in T'_{f,1}$ and an index $j \in J^w(T_{f,0})$ such that $y_j \neq w_j$, then output reject, otherwise output accept.

Figure 4: The testing algorithm for general monomials.

D). For each point y in the sample that belongs to $f^{-1}(0)$, the algorithm performs a binary search on the pair w, y . If any search fails, then the algorithm rejects (recall that in such a case it has evidence that f is not a monomial). Otherwise, for each point y in the sample that belongs to $f^{-1}(0)$, the algorithm has an index, $j^w(y) \in \Delta(y, w)$, such that $f(w^{-j^w(y)}) = 0$. Let the subset of all these indices be denoted by J . Note that by the construction of J , if f is a monomial, then for every $j \in J$, if $w_j = 1$, then the variable x_j must belong to the conjunction defining f and if $w_j = 0$, then \bar{x}_j must belong to the conjunction.

The algorithm then takes an additional sample, also of size $\Theta(\sqrt{n}/\epsilon)$, and checks whether there exists a point y in the sample that belongs to $f^{-1}(1)$ and an index $j \in J$ such that $y_j \neq w_j$. In such a case the algorithm has evidence that f is not a monomial. Viewing this in terms of the hypergraph H_f , we have that $f(w^{-j}) = 0$, $f(y) = f(w) = 1$, and both $Z(\{y, w\}) \subseteq Z(w^{-j})$ and $O(\{y, w\}) \subseteq O(w^{-j})$, so that $\{w^{-j}, y, w\} \in E(H_f)$. The pseudo-code of the algorithm is given in Figure 4.

4.3 The Analysis of the Testing Algorithm for General Monomials

We start by establishing a very simple claim. This claim explains why the algorithm can accept f in case it sees no point in the first sample that belongs to $f^{-1}(1)$.

Claim 8 *If $D(f^{-1}(1)) \leq \epsilon$ (that is, f is ϵ -close to the all-0 function), then $\text{dist}_D(f, \text{MON}) \leq \epsilon$.*

Proof: For any choice of $y \in \{0, 1\}^n$, we define the function f_y as follows: $f_y(y) = 1$ and $f_y(w) = 0$ for all $w \neq y$. Thus, for any fixed choice of y , the function f_y is a monomial (over n literals). If $f(w) = 0$ for all w , then let y be the point with minimum weight in D (so that $D(y) \leq 2^{-n}$). Otherwise, let y be any point such that $f(y) = 1$. Based on the assumption that $\epsilon > 2^{-n}$, in both cases $\text{dist}_D(f, f_y) \leq \epsilon$. Since f_y is a monomial, the proof is completed. ■

Since the initial sample S consists of $\Theta(1/\epsilon)$ points (that are generated independently according to D), we get the next corollary.

Corollary 9 *If $\text{dist}_D(f, \mathcal{MON}) > \epsilon$, then the probability that Algorithm 4 accepts because $S_{f,1}$ is empty is $\exp(-c)$ where c is the constant in the $\Theta(\cdot)$ notation for S .*

We next modify Definition 5.

Definition 7 (Empty points and representative indices (for general monomials)) *For a point $y \in f^{-1}(0)$ and a point $w \in f^{-1}(1)$, we say that y is empty with respect to w (and f) if the binary search procedure (Algorithm 3) fails when given w and y as input. We denote the set of empty points with respect to w (and f) by $Y_\emptyset^w(f)$. If y is not empty with respect to w (and f), then we let $j^w(y) \in \Delta(y, w)$ denote the index that the binary search procedure returns. We refer to this index as the representative index for y with respect to w . If $y \in Y_\emptyset^w(f)$, then we define $j^w(y)$ to be 0.*

Using the notation introduced in Algorithm 4, $J^w(T_{f,0}) = \{j^w(y) : y \in T_{f,0} \setminus Y_\emptyset^w(f)\}$. For a subset of indices $J \subseteq [n]$, let $Y_{f,1}^w(J)$ denote the set of points $y \in f^{-1}(1)$ such that $y_j \neq w_j$ for some $j \in J$, and let $Y_{f,0}^w(J)$ denote the set of points $y \in f^{-1}(0)$ for which $j^w(y) \in J$. Lemma 3 and Claim 4 are adapted as follows.

Lemma 10 *Suppose $\text{dist}_D(f, \mathcal{MON}) > \epsilon$ and consider any fixed point $w \in f^{-1}(1)$ and a sample T of $c_1\sqrt{n}/\epsilon$ points generated independently according to D . For a sufficiently large constant c_1 , with probability at least $5/6$ over the choice of T , either $T_{f,0}$ contains an empty point with respect to w (and f) or $D(Y_{f,1}^w(J(T_{f,0}))) \geq \frac{\epsilon}{4\sqrt{n}}$.*

Claim 11 *Let I be any fixed subset of $[n]$, let w be any fixed point in $f^{-1}(1)$, and consider a sample T of $s = c_1\sqrt{n}/\epsilon$ points generated independently according to D . For a sufficiently large constant c_1 , with probability at least $9/10$ over the choice of T , either $|J^w(T_{f,0}) \setminus I| \geq \sqrt{n}$ or $D(Y_{f,0}^w([n] \setminus (J^w(T_{f,0}) \cup I))) \leq \epsilon/2$.*

The proof of Claim 11 is exactly the same as the proof of Claim 4 except that each instance of $J(T_{f,0})$ needs to be replaced by $J^w(T_{f,0})$, and each instance of the form $Y_{f,0}(\cdot)$ needs to be replaced by $Y_{f,0}^w(\cdot)$. The proof of Lemma 10 is also essentially the same as the proof of Lemma 3. However, there is one subtle difference (due to the difference in the definitions of the violation hypergraphs for monotone and general monomials) and hence we give a proof sketch that includes the details of the difference.

Proof sketch of Lemma 10: The subset C is constructed in the same manner as in the proof of Lemma 3 except that $Y_\emptyset(f)$ is replaced by $Y_\emptyset^w(f)$, each instance of $Y_{f,1}(\cdot)$ is replaced by $Y_{f,1}^w(\cdot)$, the application of Claim 4 is replaced by an application of Claim 11 (and $J(T_{f,0})$ and $Y_{f,0}(\cdot)$ are replaced by $J^w(T_{f,0})$ and $Y_{f,0}^w(\cdot)$, respectively, as in the proof of Claim 11). The argument bounding the weight of C is also the same as in the proof of Lemma 3, and so it only remains to give the details establishing that C is indeed a vertex cover of H_f (as defined in Definition 6).

Consider any point $y \in f^{-1}(0)$ that is contained in at least one edge in H_f . If $y \in Y_\emptyset^w(f)$, then it is added to the cover in the first stage. Otherwise, y has a representative index $j^w(y) \in \Delta(y, w)$. Consider the iterations in the second stage of the construction of C . If for some iteration ℓ we have that $j^w(y) \in J^w(T_{f,0}^\ell)$, then the cover contains all points in $Y_{f,1}^w(\{j^w(y)\}) \subseteq Y_{f,1}^w(J^w(T_{f,0}^\ell))$. By the definition of H_f , each edge in H_f that contains y must contain some $y' \in f^{-1}(1)$ such that $y'_{j^w(y)} = y_{j^w(y)}$ (otherwise, for some edge B , it would not be true that $Z(B \setminus \{y\}) \subseteq Z(y)$ and $O(B \setminus \{y\}) \subseteq O(y)$). But for each such y' , since $y_{j^w(y)} \neq w_{j^w(y)}$, we have that $y'_{j^w(y)} \neq w_{j^w(y)}$, and so $y' \in Y_{f,1}^w(\{j^w(y)\})$. Therefore, all edges containing

y are covered after iteration ℓ . On the other hand, if $j^w(y) \notin J^w(T_{f,0}^\ell)$ for every iteration ℓ , then y is added to C in the third stage (where the argument is as in the proof of Lemma 3). \square

Lemma 10 together with Corollary 9 imply the next theorem.

Theorem 12 *Algorithm 4 is a distribution-free 1-sided-error testing algorithm for (membership in) \mathcal{MON} . Its query complexity is $O(\sqrt{n} \log n/\epsilon)$.*

The proof of Theorem 12 is essentially the same as the proof of Theorem 5, where the application of Lemma 3 is replaced by an application of Lemma 10, and the probability that the algorithm erroneously accepts due to $S_{1,f} = \emptyset$ is taken into account (using Corollary 9).

4.4 Disallowing the all-1 function

Let \mathcal{MON}' denote the class of monomials that are a conjunction of at least one literal, so that $\mathcal{MON}' = \mathcal{MON} \setminus \{\mathbf{1}\}$ (where $\mathbf{1}$ is the all-1 function). It is possible to augment our testing algorithm for \mathcal{MON} (Algorithm 4) so that it works for \mathcal{MON}' in a very similar manner to what was shown for monotone monomials in Subsection 3.4. The only difference is that when we take the additional sample R , we check whether there exists at least one index $i \in [n]$ such that for some $b \in \{0, 1\}$, $y_i = b$ for all points $y \in S \cap f^{-1}(1)$. The modification in the analysis is that we show that if $\text{dist}_D(f, \mathbf{1}) \leq \epsilon/2$ and $\text{dist}_D(f, \mathcal{MON}') > \epsilon$, then $\Pr_{y \sim D}[f(y) = 1 \text{ and } y_i = b] \geq \epsilon/2$ for every $i \in [n]$ and $b \in \{0, 1\}$.

Acknowledgements

We would like to thank an anonymous reviewer of STOC 2010 for suggesting to simplify the proof of Lemma 2, and to thank the anonymous reviewers of RANDOM 2010 for helpful comments.

References

- [BLR93] M. Blum, M. Luby, and R. Rubinfeld. Self-testing/correcting with applications to numerical problems. *JACM*, 47:549–595, 1993.
- [DGL⁺99] Y. Dodis, O. Goldreich, E. Lehman, S. Raskhodnikova, D. Ron, and A. Samorodnitsky. Improved testing algorithms for monotonicity. In *Proceedings of the 3rd RANDOM*, pages 97–108, 1999.
- [DLM⁺07] I. Diakonikolas, H. K. Lee, K. Matulef, K. Onak, R. Rubinfeld, R. A. Servedio, and A. Wan. Testing for concise representations. In *Proceedings of the 48th FOCS*, pages 549–557, 2007.
- [EKK⁺00] F. Ergun, S. Kannan, S. R. Kumar, R. Rubinfeld, and M. Viswanathan. Spot-checkers. *JCSS*, 60(3):717–751, 2000.
- [FKR⁺04] E. Fischer, G. Kindler, D. Ron, S. Safra, and S. Samorodnitsky. Testing juntas. *JCSS*, 68(4):753–787, 2004.
- [GGL⁺00] O. Goldreich, S. Goldwasser, E. Lehman, D. Ron, and A. Samordnitsky. Testing monotonicity. *Combinatorica*, 20(3):301–337, 2000.

- [GGR98] O. Goldreich, S. Goldwasser, and D. Ron. Property testing and its connection to learning and approximation. *JACM*, 45(4):653–750, 1998.
- [GS07] D. Glasner and R. A. Servedio. Distribution-free testing lower bounds for basic Boolean functions. In *Proceedings of the 11th RANDOM*, pages 494–508, 2007.
- [HK03] S. Halevy and E. Kushilevitz. Distribution-free property testing. In *Proceedings of the 7th RANDOM*, pages 341–353, 2003.
- [HK04] S. Halevy and E. Kushilevitz. Distribution-free connectivity testing. In *Proceedings of the 8th RANDOM*, pages 393–404, 2004.
- [HK05] S. Halevy and E. Kushilevitz. A lower bound for distribution-free monotonicity testing. In *Proceedings of the 9th RANDOM*, pages 330–341, 2005.
- [HK07] S. Halevy and E. Kushilevitz. Distribution-free property testing. *SICOMP*, 37(4):1107–1138, 2007.
- [KS09] S. Kopparty and S. Saraf. Tolerant linearity testing and locally testable codes. In *Proceedings of the 14th RANDOM*, pages 601–614, 2009.
- [MORS09] K. Matulef, R. O’Donnell, R. Rubinfeld, and R. A. Servedio. Testing halfspaces. In *Proceedings of the 20th SODA*, pages 256–264, 2009.
- [PRR06] M. Parnas, D. Ron, and R. Rubinfeld. Tolerant property testing and distance approximation. *JCSS*, 72(6):1012–1042, 2006.
- [PRS02] M. Parnas, D. Ron, and A. Samorodnitsky. Testing basic boolean formulae. *SIDMA*, 16(1):20–46, 2002.
- [RS96] R. Rubinfeld and M. Sudan. Robust characterization of polynomials with applications to program testing. *SICOMP*, 25(2):252–271, 1996.
- [Tur93] G. Turán. Lower bounds for PAC learning with queries. In *Proceedings of the 6th COLT*, pages 384–391, 1993.
- [Val84] L. G. Valiant. A theory of the learnable. *CACM*, 27(11):1134–1142, November 1984.